



Molecular Pharmacology at UPV

Molecular Pharmacology Blog by students of UPV Biotechnology degree

About



- Joserra Murguia
-

[Subscribe by email »](#)

Get the latest updates in your email box automatically.

[Subscribe via RSS »](#)

Scientists identify the largest network of protein interactions related to Alzheimers disease

Alzheimer's disease is an age-related neurodegenerative disease. Despite the considerable efforts made in recent years to understand the mechanisms that trigger this disease, an effective treatment is not yet available. This study reveals new molecular and functional data that could help researchers gain a better understanding of the disease and potentially to develop new therapies.

Proteins are the molecular instruments that cells use to carry out their functions. Proteins don't normally act alone, but interact with other proteins to form cellular networks. In this study, rather than looking at individual proteins as been done in many previous studies, the scientists analyzed the biology behind Alzheimer's disease using a global approach.

To do so, the researchers used genetic methods to study the thousands of possible interactions between proteins known or thought likely - to be involved in the disease, including proteins derived from the genes located on the chromosomes related to the Alzheimer's disease. They obtained a total of 200 new interactions. This information, added to what is already known, brings the total number of Alzheimer's-related interactions to 6000, and involves 1700 proteins, resulting in the largest network of interactions between proteins related to Alzheimer's disease.

The computational analysis of these interactions, performed by the MareNostrum Supercomputer of the Barcelona Supercomputing Center (BSC), revealed that many of the groups of proteins are highly interconnected and have similar functions in cells. Some of these functions suggest new molecular mechanisms that could be linked to Alzheimer's. A clear example is a protein called ECSIT, that relates oxidative stress to inflammation and changes in the mitochondria, indicating that these processes may play a role in the development of the disease.

Increased life expectancy and improvements in health care and diagnostics have led to the increased prevalence of the Alzheimer's disease, which is a serious problem in both developed and developing countries. This study opens the door to new research aimed at finding a cure for the disease.

Reference article:

Interactome mapping suggests new mechanistic details underlying Alzheimer's disease

Montserrat Soler-López, Andreas Zanzoni, Ricart Lluís, Ulrich Stelzl and Patrick Aloy

Genome Research (2010) [DOI: 10.1101/gr.114280.110]

Comments (0)

Add a Comment


To leave a comment on this posterous, please login by clicking one of the following.



Meta

- Posted at 1:17 pm on Thu, Dec 30, 2010
- Views (24) [Permalink](#) (∞)
- [Comments \(0\)](#) [Add A Comment](#)

Share

- [Short URL](#)
-
- [Headline by Airspace Workshop for Posterous](#)
-  **3704** site views

[Create your own free Posterous site now »](#)